

# THE GROWTH OF GRIGORCHUK'S TORSION GROUP

LAURENT BARTHOLDI

ABSTRACT. In 1980 Rostislav Grigorchuk constructed a group  $G$  of intermediate growth, and later obtained the following estimates on its growth  $\gamma$  [Gri84]:

$$e^{\sqrt{n}} \lesssim \gamma(n) \lesssim e^{n^\beta},$$

where  $\beta = \log_{32}(31) \approx 0.991$ . Using elementary methods we improve the upper bound to

$$\gamma(n) \lesssim e^{n^\alpha},$$

where  $\eta \approx 0.811$  is the real root of the polynomial  $X^3 + X^2 + X - 2$  and  $\alpha = \log(2)/\log(2/\eta) \approx 0.767$ .

## 1. INTRODUCTION

The notion of growth for finitely generated groups was introduced in the 1950's in the former USSR [Sva55] and in the 1960's in the West [Mil68]. There are well-known classes of groups of polynomial growth (abelian, and more generally virtually nilpotent groups [Gro81]) and of exponential growth (non-virtually-nilpotent linear [Tit72] or non-elementary hyperbolic [GH90] groups). However, the first example of a group of intermediate growth was discovered later, by Rostislav Grigorchuk; see [Gri83, Gri84, Gri91]. He showed that the growth  $\gamma$  of his group satisfies

$$e^{\sqrt{n}} \lesssim \gamma(n) \lesssim e^{n^\beta},$$

where  $\beta = \log_{32}(31) \approx 0.991$ ; see below for the precise definition of growth. The purpose of this note is to prove the following improvement:

**Theorem 1.1.** *Let  $\eta$  be the real root of the polynomial  $X^3 + X^2 + X - 2$ , and set  $\alpha = \log(2)/\log(2/\eta) \approx 0.767$ . Then the growth  $\gamma$  of Grigorchuk's group satisfies*

$$e^{\sqrt{n}} \lesssim \gamma(n) \lesssim e^{n^\alpha}.$$

## 2. GROWTH OF GROUPS

Let  $G$  be a group generated as a monoid by a finite set  $S$ . A *weight* on  $(G, S)$  is a function  $\omega : S \rightarrow \mathbb{R}_+$ . It induces a *length*  $\partial_\omega$  on  $G$  by

$$\partial_\omega : \begin{cases} G \rightarrow \mathbb{R}_+ \\ g \mapsto \min\{\omega(s_1) + \cdots + \omega(s_n) \mid s_1 \cdots s_n =_G g\}. \end{cases}$$

A *minimal form* of  $g \in G$  is a representation of  $g$  as a word of minimal length over  $S$ . The *growth* of  $G$  with respect to  $\omega$  is then

$$\gamma_\omega : \begin{cases} \mathbb{R}_+ \rightarrow \mathbb{R}_+ \\ n \mapsto \#\{g \in G \mid \partial_\omega(g) \leq n\}. \end{cases}$$

The function  $\gamma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is *dominated* by  $\delta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ , written  $\gamma \lesssim \delta$ , if there is a constant  $C \in \mathbb{R}_+$  such that  $\gamma(n) \leq \delta(Cn)$  for all  $n \in \mathbb{R}_+$ . Two functions  $\gamma, \delta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  are *equivalent*, written  $\gamma \sim \delta$ , if  $\gamma \lesssim \delta$  and  $\delta \lesssim \gamma$ .

The following lemmata are well known:

---

*Date:* February 1, 2008.

This work has been supported by the "Swiss National Fund for Scientific Research".

**Lemma 2.1.** *Let  $S$  and  $S'$  be two finite generating sets for the group  $G$ , and let  $\omega$  and  $\omega'$  be weights on  $(G, S)$  and  $(G, S')$  respectively. Then  $\gamma_\omega \sim \gamma_{\omega'}$ .*

*Proof.* Let  $C = \max_{s \in S} \partial_{\omega'}(s)/\omega(s)$ . Then  $\partial_{\omega'}(g) \leq C\partial_\omega(g)$  for all  $g \in G$ , and thus  $\gamma_\omega(n) \leq \gamma_{\omega'}(Cn)$ , from which  $\gamma_\omega \lesssim \gamma_{\omega'}$ . The opposite relation holds by symmetry.  $\square$

The *growth type* of a finitely generated group  $G$  is the  $\sim$ -equivalence class containing its growth functions; it will be denoted by  $\gamma_G$ .

Note that all exponential functions  $b^n$  are equivalent, and polynomial functions of different degree are inequivalent; the same holds for the subexponential functions  $e^{n^\alpha}$ . We have

$$0 \not\lesssim n \not\lesssim n^2 \cdots \not\lesssim e^{\sqrt{n}} \not\lesssim \cdots \not\lesssim e^n.$$

Note also that the ordering  $\lesssim$  is not linear.

**Lemma 2.2.** *Let  $G$  be a finitely generated group. Then  $\gamma_G \lesssim e^n$ .*

*Proof.* Choose for  $G$  a finite generating set  $S$ , and define the weight  $\omega$  by  $\omega(s) = 1$  for all  $s \in S$ . Then  $\gamma_\omega(n) \leq |S|^n$  for all  $n$ , so  $\gamma_G \lesssim e^n$ .  $\square$

If there is a  $d \in \mathbb{N}$  such that  $\gamma_G \lesssim n^d$ , the group  $G$  is of *polynomial growth* of degree at most  $d$ ; if  $\gamma_G \sim e^n$ , then  $G$  is of *exponential growth*; otherwise  $G$  is of *intermediate growth*. The existence of groups of intermediate growth was first shown by Grigorchuk [Gri83].

### 3. THE GRIGORCHUK 2-GROUP

Let  $\Sigma^*$  be the set of finite sequences over  $\Sigma = \{0, 1\}$ . For  $x \in \Sigma$  set  $\bar{x} = 1 - x$ . Define recursively the following length-preserving permutations of  $\Sigma^*$ :

$$\begin{aligned} a(x\sigma) &= \bar{x}\sigma; \\ b(0\sigma) &= 0a(\sigma), & b(1\sigma) &= 1c(\sigma); \\ c(0\sigma) &= 0a(\sigma), & c(1\sigma) &= 1d(\sigma); \\ d(0\sigma) &= 0\sigma, & d(1\sigma) &= 1b(\sigma). \end{aligned}$$

Then  $G$ , the Grigorchuk 2-group [Gri80, Gri84], is the group generated by  $S = \{a, b, c, d\}$ . It is readily checked that these generators are of order 2 and that  $V = \{1, b, c, d\}$  is a Klein group.

Let  $H = V^G$  be the normal closure of  $V$  in  $G$ . It is of index 2 in  $G$  and preserves the first letter of sequences; i.e.  $H \cdot x\Sigma^* \subset x\Sigma^*$  for all  $x \in \Sigma$ . There is a map  $\psi : H \rightarrow G \times G$ , written  $g \mapsto (g_0, g_1)$ , defined by  $0g_0(\sigma) = g(0\sigma)$  and  $1g_1(\sigma) = g(1\sigma)$ . As  $H = \langle b, c, d, b^a, c^a, d^a \rangle$ , we can write  $\psi$  explicitly as

$$\psi : \begin{cases} b \mapsto (a, c), & b^a \mapsto (c, a) \\ c \mapsto (a, d), & c^a \mapsto (d, a) \\ d \mapsto (1, b), & d^a \mapsto (b, 1). \end{cases}$$

### 4. THE GROWTH OF $G$

Let  $\eta \approx 0.811$  be the real root of the polynomial  $X^3 + X^2 + X - 2$ , and define the following function on  $S$ :

$$\begin{aligned} \omega(a) &= 1 - \eta^3 = \eta^2 + \eta - 1, & \omega(c) &= 1 - \eta^2, \\ \omega(b) &= \eta^3 = 2 - \eta - \eta^2, & \omega(d) &= 1 - \eta. \end{aligned}$$

It is a weight, because it takes positive values on every generator.

**Lemma 4.1.** *Every  $g \in G$  admits a minimal form*

$$[*]a * a * a \cdots * a[*],$$

where  $*$   $\in \{b, c, d\}$  and the first and last  $*$ s are optional.

*Proof.* Clearly  $\omega(s) > 0$  for  $s \in S$ , so  $\omega$  is a weight. Let  $w$  be a minimal form of  $g$ . The lemma asserts that one can suppose there are no consecutive letters in  $\{b, c, d\}$  in  $w$ ; now two equal consecutive letters cancel, and the product of any two distinct letters in  $\{b, c, d\}$  equals the third one. For any arrangement  $(x, y, z)$  of  $\{b, c, d\}$  we have  $\omega(x) \leq \omega(y) + \omega(z)$ , so the substitution of  $z$  for  $xy$  will not increase the weight of  $w$ .  $\square$

**Proposition 4.2.** *Let  $g \in H$ , with  $\psi(g) = (g_0, g_1)$ . Then*

$$\eta(\partial_\omega(g) + \omega(a)) \geq \partial_\omega(g_0) + \partial_\omega(g_1).$$

*Proof.* Let  $w$  be a minimal form of  $g$ . Thanks to Lemma 4.1 we may suppose the number of  $*$ s in  $w$  is at most the number of  $a$ s plus one. Construct words  $w_0, w_1$  over  $S$  using  $\psi$  seen as a substitution on words; they represent  $g_0$  and  $g_1$  respectively. Note that

$$\begin{aligned} \eta(\omega(a) + \omega(b)) &= \omega(a) + \omega(c), \\ \eta(\omega(a) + \omega(c)) &= \omega(a) + \omega(d), \\ \eta(\omega(a) + \omega(d)) &= 0 + \omega(b). \end{aligned}$$

As  $\psi(b) = (a, c)$  and  $\phi(aba) = (c, a)$ , each  $b$  in  $w$  contributes  $\omega(a) + \omega(c)$  to the total weight of  $w_0$  and  $w_1$ ; the same argument applies to  $c$  and  $d$ . Now, grouping together pairs of  $*$ s in  $\{b, c, d\}$  and  $as$ , we see that  $\eta(\partial_\omega(g))$  is a sum of left-hand terms, possibly  $-\eta\omega(a)$ ; while  $\partial_\omega(g_0) + \partial_\omega(g_1)$  is bounded by the total weight of the letters in  $w_0$  and  $w_1$ , which is precisely the sum of the corresponding right-hand terms.  $\square$

Let  $\alpha = \log(2)/\log(2/\eta) \approx 0.767$ , and for  $n \in \mathbb{N}$  set  $C_n = \frac{1}{n+1} \binom{2n}{n}$ , the  $n$ th Catalan number; remember that it is the number of labelled binary rooted trees with  $n+1$  leaves [Cat38, LW92, page 119].

**Proposition 4.3.** *Let  $\zeta = \frac{\omega(a)}{2/\eta-1}$ , let  $K > \zeta$  be any constant, and for  $n \in \mathbb{R}_+$  set*

$$L_n = \max \left\{ 1, \left\lceil 2 \left( \frac{n-\zeta}{K-\zeta} \right)^\alpha \right\rceil - 1 \right\}.$$

*Then we have*

$$(1) \quad \gamma_\omega(n) \leq C_{L_n-1} 2^{L_n-1} \gamma_\omega(K)^{L_n}.$$

*Proof.* We construct an injection  $\iota$  of  $G$  into the set of labelled binary rooted trees each of whose leaves is labelled by an element of  $G$  of weight bounded by  $K$  and each of whose interior vertices is labelled by an element of the subgroup  $\langle a \rangle$  of  $G$ . For  $g \in G$ ,  $\iota(g)$  is called its *representation*. It is constructed as follows: if  $g \in G$  satisfies  $\partial_\omega(g) \leq K$ , its representation is a tree with one vertex labelled by  $g$ . If  $\partial_\omega(g) > K$ , let  $h \in \langle a \rangle$  be such that  $gh \in H$ , and write  $\psi(gh) = (g_0, g_1)$ . By Proposition 4.2,  $\partial_\omega(g_i) \leq \eta \partial_\omega(g)$ , so we may construct inductively the representations of  $g_0$  and  $g_1$ . The representation of  $g$  is a tree with  $h$  at its root vertex and  $\iota(g_0)$  and  $\iota(g_1)$  attached to its two branches.

We first claim that  $\iota$  is injective: let  $\mathcal{T}$  be a tree in the image of  $\iota$ . If  $\mathcal{T}$  has one node labelled by  $g$ , then  $\iota^{-1}(\mathcal{T}) = \{g\}$ . If  $\mathcal{T}$  has more than one vertex, let  $h \in \langle a \rangle$  be the label of the root vertex and  $(\mathcal{T}_0, \mathcal{T}_1)$  be the two subtrees connected to the root vertex. By induction on the number of vertices of  $\mathcal{T}$ , we have  $\mathcal{T}_i = \iota(g_i)$  for unique  $g_0$  and  $g_1$ . Then as  $\psi$  is injective there is a unique  $g \in G$  with  $\psi(gh) = (g_0, g_1)$ , and  $\iota^{-1}(\mathcal{T}) = \{g\}$ .

We next prove by induction on  $n$  that if  $\partial_\omega(g) \leq n$  then its representation is a tree with at most  $L_n$  leaves. Indeed if  $n \leq K$  then  $g$ 's representation has one leaf and  $L_n = 1$ , while otherwise  $g$ 's representation is made up of those of  $g_0$  and  $g_1$ . Say  $\partial_\omega(g_0) = \ell$  and  $\partial_\omega(g_1) = m$ ; then by Proposition 4.2 we have  $\ell + m \leq \eta(n + \omega(a))$ . By induction these representations have at most  $L_\ell$  and  $L_m$  leaves. As  $\alpha < 1$ , we have  $L_\ell + L_m \leq 2L_{(\ell+m)/2}$  for all  $\ell, m$ ; and by direct computation,  $L_{\eta/2(n+\omega(a))} = \lfloor L_n/2 \rfloor$ , so the number of leaves of  $g$ 's representation is

$$L_\ell + L_m \leq 2L_{(\ell+m)/2} \leq 2L_{\eta/2(n+\omega(a))} \leq L_n,$$

as was claimed.

We conclude that  $\gamma(n)$  is bounded by the number of representations with  $L_n$  leaves; there are  $C_{L_n-1}$  binary trees with  $L_n$  leaves, 2 choices of labelling for each of the  $L_n - 1$  interior vertices, and  $\gamma(K)$  choices for each leaf; so Equation (1) follows.  $\square$

A lower bound on the growth of  $G$  comes from the fact that  $G$  is residually a 2-group:

**Theorem 4.4** (Grigorchuk [Gri89]). *Suppose  $G$  is a finitely generated residually- $p$  group. Let  $\{G_n\}$  be the Zassenhaus filtration of  $G$ . If  $[G : G_n] \gtrsim n^d$  for all  $d$ , then  $\gamma_G \gtrsim e^{\sqrt{n}}$ .*

*Proof of Theorem 1.1.* The sequence  $[G : G_n]$  was shown to be of superpolynomial growth in [Gri89], so Theorem 4.4 yields the claimed lower bound; an elementary proof of this lower bound appears also in [Gri84].

For the upper bound, which is the main result of the present note, we invoke Proposition 4.3 with  $K = 1$ , noting that  $L_n \sim n^\alpha$  and  $C_{L_n} \leq 4^{L_n}$ , to obtain  $\gamma_\omega \lesssim (4 \cdot 2 \cdot \gamma(1))^{n^\alpha} \sim e^{n^\alpha}$ .  $\square$

## 5. CONCLUSION

The main fact used in the proof of Theorem 1.1 is the existence of minimal forms given by Lemma 4.1, coming from the natural map  $\langle a \rangle * V \twoheadrightarrow G$ . One can impose stronger conditions on minimal forms, such as ‘not containing *dada* as a subword’, coming from an explicit recursive presentation of  $G$  [Lys85]. Tighter upper bounds result from such considerations. Yuriy Leonov [Leo98] recently obtained improvements on the lower bound of Theorem 1.1.

I wish to thank Robyn Curtis, Igor Lysënok and Pierre de la Harpe for having patiently listened to preliminary—and incorrect—proofs of these results, and also of course Rostislav Grigorchuk without whom these results couldn’t even have existed.

## REFERENCES

- [Cat38] Eugène C. Catalan, *Note sur une équation aux différences finies*, J. Math. Pures Appl. (9) **3** (1838), 508–516.
- [GH90] Étienne Ghys and Pierre de la Harpe, *Sur les groupes hyperboliques d’après Mikhael Gromov*, Progress in Mathematics, vol. 83, Birkhäuser Boston Inc., Boston, MA, 1990, Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988.
- [Gri80] Rostislav I. Grigorchuk, *On Burnside’s problem on periodic groups*, Funktsional. Anal. i Prilozhen. **14** (1980), no. 1, 53–54, English translation: Functional Anal. Appl. **14** (1980), 41–43.
- [Gri83] Rostislav I. Grigorchuk, *On the Milnor problem of group growth*, Dokl. Akad. Nauk SSSR **271** (1983), no. 1, 30–33.
- [Gri84] Rostislav I. Grigorchuk, *Degrees of growth of finitely generated groups and the theory of invariant means*, Izv. Akad. Nauk SSSR Ser. Mat. **48** (1984), no. 5, 939–985, English translation: Math. USSR-Izv. **25** (1985), no. 2, 259–300.
- [Gri89] Rostislav I. Grigorchuk, *On the Hilbert-Poincaré series of graded algebras that are associated with groups*, Mat. Sb. **180** (1989), no. 2, 207–225, 304, English translation: Math. USSR-Sb. **66** (1990), no. 1, 211–229.
- [Gri91] Rostislav I. Grigorchuk, *On growth in group theory*, Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990) (Tokyo), Math. Soc. Japan, 1991, pp. 325–338.
- [Gro81] Mikhael Gromov, *Groups of polynomial growth and expanding maps*, Inst. Hautes Études Sci. Publ. Math. (1981), no. 53, 53–73.
- [Leo98] Yuriy G. Leonov, *On lower estimation of growth for some torsion groups*, to appear, 1998.
- [LW92] Jacobus H. van Lint and Richard M. Wilson, *A course in combinatorics*, Cambridge University Press, 1992.
- [Lys85] Igor G. Lysionok, *A system of defining relations for the Grigorchuk group*, Mat. Zametki **38** (1985), 503–511.
- [Mil68] John W. Milnor, *Growth of finitely generated solvable groups*, J. Differential Geom. **2** (1968), 447–449.
- [Sva55] A. S. Svarts, *A volume invariant of coverings*, Dokl. Akad. Nauk SSSR (1955), no. 105, 32–34 (Russian).
- [Tit72] Jacques Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270.

SECTION DE MATHÉMATIQUES  
UNIVERSITÉ DE GENÈVE  
CP 240, 1211 GENÈVE 24  
SWITZERLAND